



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Virology

CrossMark

The emergence and cross species transmission of newly discovered tick-borne Bunyavirus in China

Yong-Zhen Zhang and Jianguo Xu

A novel tick-borne Bunyavirus, discovered in China and later in South Korea and Japan, is now known as Huaiyangshan virus or severe fever with thrombocytopenia syndrome virus and has been identified as the causative agent of a hemorrhagic fever-like disease. Of five species of ticks carrying Huaiyangshan viruses, *Haemaphysalis longicornis* was the most abundant in regions where the virus was endemic. Its usual hosts (cattle, goats, dogs, rats and chickens) tested positive for Huaiyangshan virus RNA and had a high seroprevalence. The distribution of *H. longicornis* and the migratory routes of four wild fowl across China, South Korea and Japan are coincident. Thus a tick and migratory bird model for the transmission of the Huaiyangshan virus was proposed.

Address

State Key Laboratory for Infectious Disease Prevention and Control, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Changping, Beijing 102206, China

Corresponding author: Xu, Jianguo (xujianguo@icdc.cn)

Current Opinion in Virology 2016, **16**:126–131

This review comes from a themed issue on **Emerging viruses: interspecies transmission**

Edited by **Christian Drosten** and **Yi Guan**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 4th March 2016

<http://dx.doi.org/10.1016/j.coviro.2016.02.006>

1879-6257/© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Introduction

In 2009, a novel hemorrhagic fever-like disease was first observed in China [1] and a novel Bunyavirus was identified as the causative agent in 2010 [1]. Surveillance and epidemiologic investigations confirmed its wide geographic distribution in China [2]. From 2010 to 2013, 2047 cases of human infections were reported, and the virus was found to be present (designated as endemic) in 16 provinces in the central and eastern regions in China [3]. 94.4% of these cases were farmers or forest workers, who lived in rural areas and engaged in the agricultural activities [3,4]. Subsequently, the disease was identified in South Korea and Japan in 2012 [5–7]. Retrospective studies dated the earliest known human infections back to 2005 in Japan and 2006 in China [6,8]. The broad

geographic distribution of the virus across China, Korea and Japan, its high fatality rate, and the potential for human-to-human transmission led this virus to be seen as an increasingly important threat to regional and global health [9].

Emergence of the new virus

In April 2009, physicians of the Union Hospital of Tongji Medical College, Wuhan, China noticed several fatalities among a cluster of patients with high fever, leucopenia, thrombocytopenia, and elevated liver transaminase levels [1,10,11^{*}]. All patients were from the Huaiyangshan mountain area of Hubei and Henan provinces, in the central part of China. Some of the patients recalled a history of tick bites [1,10,11^{*}]. As the clinical features were highly similar to those of human granulocytic anaplasmosis (HGA), caused by the bacterium *Anaplasma phagocytophilum* and first reported in the neighboring province of Anhui in 2006 [12], the illnesses were clinically diagnosed as suspected HGA in 2010 [1,10,11^{*}]. Samples from patients were sent to the National Institute for Communicable Disease Control and Prevention of the China CDC for confirmation of HGA infection. HGA was immediately excluded as the cause since all specimens tested negative for *A. phagocytophilum* by PCR and immunologic methods [12]. Fortunately, attempts to isolate the etiologic agent by culturing in DH82 cells were finally successful, leading to the discovery of a novel Bunyavirus, which was published in March 2011 in Chinese Journal of Epidemiology [1].

As several groups were studying the disease and its agent in the initial phase, it was reported as Huaiyangshan hemorrhagic fever (HYSHF) [1], severe fever with thrombocytopenia syndrome (SFTS) [2], and fever, thrombocytopenia and leukopenia syndrome (FTLS) [13]. Accordingly, the etiologic agent was named as Huaiyangshan virus (HYSV), severe fever with thrombocytopenia syndrome virus (SFTSV) and Henan fever virus (HFV). As the geographic origin of the initially reported patients was the Huaiyangshan mountain area and severe fever and thrombocytopenia are not specific for any given disease, but present in nearly all hemorrhagic fevers caused by other viruses, we recommend that the disease be referred to as Huaiyangshan hemorrhagic fever (HYSHF) and the causative virus as Huaiyangshan virus (HYSV) [1,14]. These were the names used in the first published report on this disease and virus, although it appeared in a Chinese language journal [1], and are used in this review. To date, human infections with HYSV

have also been reported in South Korea and Japan [5,6,15].

Genetic characteristic of HYSV

Huaiyangshan virus belongs to the genus *Phlebovirus* in the family *Bunyaviridae* and has been isolated successfully from human and animal samples in China. Like other members of the *Bunyaviridae*, HYSV is an enveloped, negative-stranded RNA virus [1]. Its genome consists of three RNA segments: small (S), medium (M), and large (L). The L segment has a length of 6,368 nucleotides, with an 11 nucleotide 5'-noncoding region (NCR), an open reading frame (ORF) encoding the RNA-dependant RNA polymerase (RdRp) of 2084 amino acids, and a 113 nucleotide 3'-NCR [16]. The M segment has a length of 3378 nucleotides, with an 18-nucleotide 5'-NCR, an ORF coding for a Glycoprotein N and C (GnGc) precursor of 1073 amino acids, and a 138-nucleotide-long 3'-NCR. Similar to the Uukuniemi-like viruses, HYSV does not have the non-structural protein found in the Phlebotomus fever viruses, however its two surface glycoproteins are in the same order (Gn-Gc) as both these groups. The S segment has a length of 1746 nucleotides, with a 43-nucleotide 5'-NCR, an ORF coding for an N protein, an ORF coding in the opposite sense for an NSs protein and a 29-nucleotide 3'-NCR. A typical characteristic of *Bunyaviridae* family viruses is that the 3' and 5' ends of each RNA segment are conserved and form stable base-paired panhandles. In HYSV, the L and M segments have one nucleotide different from the consensus at the 5' end.

The HYSVs from China, South Korea, and Japan are closely related to each other [16,17^{••},18,19^{••}] with a nucleotide divergence of less than 6% for all three segments. To date, all known HYSVs identified in China can be classified into six lineages [14,16,18] and there are three lineages of HYSVs in Japan [19^{••}] with the third lineage from Japan closely related to the HYSVs from Zhejiang Province, China. These data suggest that these viruses have a recent common ancestor, likely due to migration of their hosts. Genetic recombination and reassortment between different HYSVs might have occurred [20,21].

Clinical presentations of HYSHF are nonspecific

The clinical presentations of HYSHF are nonspecific, and are difficult to distinguish from those of HGA, hemorrhagic fever with renal syndrome (HFRS), and leptospirosis. The first group of patients presenting to the Union Hospital of Tongji Medical College, Hubei Province in 2009, were initially suspected to have HGA, before a diagnosis of HYSHF was made [1].

Haemaphysalis longicornis ticks are the major vector

Investigation of the first 33 laboratory-confirmed cases revealed that all patients were farmers, living in villages

along the low Huaiyangshan mountain area, and involved in domestic animal herding, tea collection and other agricultural activities [2]. From April to May 2010, 613 adult ticks were collected from domestic animals around the villages where these patients lived and nearby regions. Only two species were identified; *H. longicornis* (88.09%) and *Boophilus microplus* (11.91%). The HYSV virus tested positive in the *H. longicornis* (4.93%) and *R. microplus* (0.613%) tick pools. The HYSV-positive *H. longicornis* ticks were only found in the endemic regions (where patients had been infected with HYSV), whereas HYSV-positive *R. microplus* ticks were found in both endemic and nonendemic regions. On the basis of this observation, we hypothesized that *H. longicornis* was the major vector of HYSV [1].

To test this hypothesis, we extended the sampling sites to 15 endemic counties of Hubei and Henan provinces, and 23 non-endemic counties of Hubei Province, around the Huaiyangshan area. In the 15 endemic counties, *H. longicornis* ticks ($N = 3498$) were more abundant than *R. microplus* ticks ($N = 1003$). In the 23 nonendemic counties, *R. microplus* ($N = 12,830$) was more abundant than *H. longicornis* ($N = 400$) [16]. As in the smaller study [1], the HYSV-positive *H. longicornis* ticks were found only in the endemic counties while HYSV-positive *R. microplus* ticks were found in both endemic and non-endemic counties. These observations supported the hypothesis that *H. longicornis* is the major vector of HYSV [16].

A similar situation was found in Shangdong Province of China, where infections have been identified since 2010. Wang *et al.* reported that 96.9% ticks ($N = 3145$) collected were *H. longicornis*, followed by *R. microplus* (2.6%), *H. campanulata* (0.3%) and *Dermacentor sinicus* (0.2%). HYSV was only detected in *H. longicornis* (96.3%) and adult *R. microplus* (3.7%) [22[•]]. Recently, Meng *et al.* detected HYSV in 4.17% ($n = 72$) of *H. concinna* ticks from goats [23].

In South Korea, Park *et al.* found that 90.8% of ticks ($N = 13,053$) collected from 2011 to 2012 were *H. longicornis*, followed by *H. flava* (8.8%), *Ixodes nipponensis* (0.3%) and *I. persulcatus* (0.05%). The HYSV-positive ticks were widely distributed in the country and an HYSV infection was confirmed in 2012 in Gangwon Province where virus-positive ticks had been previously detected [5,24].

Yun *et al.* collected 261 ticks from humans during May–October 2013 in South Korea. They found that the most abundant tick from humans was *H. longicornis* (81.2%), followed by *Amblyomma testudinarium* (6.5%); *I. nipponensis* (5.7%); *H. flava* (5.4%), *H. japonica*, *I. persulcatus*, and *I. granulatus* (0.4% each). The minimum infection rate per 100 ticks (MIR) was 5.7% in *H. longicornis*, 23.5% in *A. testudinarium* and 13.3% in *I. nipponensis* ticks [25^{••}]. This finding of other species of ticks harboring HYSV should

be further addressed. However, it is clear that *H. longicornis* ticks are the major species in the environment of endemic HYSV areas and are the major vector for HYSV [5,16,25[•],26].

Domestic animals are widely infected

Zhang *et al.* found that some ticks from cattle, goats, dogs and cats in the Huaiyangshan mountain areas tested positive for HYSV [1,16]. Wang *et al.* reported that the prevalence of HYSV-positive ticks from sheep and dogs was higher than that in ticks from cattle in Shandong Province [22[•]]. Ding *et al.* noted that the seroprevalence for HYSV was 41.8% (268/641) in domestic animals in the HYSV endemic region of Laizhou City, Shandong Province; specifically, goats (74.8%), cattle (57.1%), dogs (52.1%), chickens (35.9%) and pigs (0%) [27]. Niu *et al.* sampled 3039 animals in two endemic counties of Shandong Province from April to November 2011, and found that seroprevalence for HYSV was 69.5% for sheep, 60.5% for cattle, 37.9% for dogs, 3.1% for pigs, and 47.4% for chickens [28]. Li *et al.* found that in six endemic counties of Jiangsu Province from March 2012 to February 2013, the seroprevalence in goats, cattle, dogs, pigs, chickens, geese, rodents and hedgehogs was 66.8%, 28.2%, 7.4%, 4.7%, 1.2%, 1.7%, 4.4% and 2.7%, respectively [29[•]]. In all these cases serum samples were tested by double-antigen sandwich ELISA. Using the same method, Xing *et al.* found that seroprevalence ranged from 10% to 18% in samples from cattle, sheep, goats, deer, and elk in 24 Minnesota


counties of the USA [30]. A possible cross-reaction with Heartland virus, which is phylogenetically related to HYSV and extensively identified in USA, should be considered when interpreting this result [31].

HYSV RNA has also been detected in some animals. Ni *et al.* investigated rodents as possible hosts of this virus and captured 40 *Rattus norvegicus* in villages near the houses of patients as well as eight *Apodemus agrarius* and four *Rattus losea* in the mountainous regions near villages where human cases of HYSHF had been reported in Zhejiang Province from 2012 to 2014. Two of the eight *A. agrarius* rats were positive for HYSV RNA and virus was isolated from the lung tissues of the two animals. The genomic sequences of the rat viruses had a high level of sequence identity to those isolated from patients [32[•]]. Although seropositive rodents have also been reported, more study is required to clarify if rodents have a role in the transmission of the virus [29[•]]. Niu *et al.* reported that HYSV RNA was detected in all sampled animal species, ranging from 1.7% to 5.3%, and viruses were isolated from sheep, cattle, and dogs [28]. Again, the role played by these animals in transmission of the virus needs further study.

A tick and migratory bird model for HYSV transmission

Using whole genome sequences, Yun *et al.* analyzed the phylogenetic relationships among HYSVs and showed

Figure 1



Tick and migratory bird pathway for the transmission of HYSV. The thicker solid and dashed arrows from domestic animals and agricultural activity indicate that virus was isolated or viral RNA was detected, respectively. The dashed arrows in wildlife and activity indicate viral RNA was detected in ticks captured from those wild animals and grassland.

that it had been circulating between China, South Korea, and Japan and that HYSV-infected *H. longicornis* and human infections had been found in all three countries. They hypothesized that HYSV was transmitted, at least in part, by the *H. longicornis* tick and moved over long distances to Korea and Japan through migratory birds, especially across water bodies of the East China Sea, Yellow Sea, and the East Sea/Sea of Japan [33**].

This hypothesis is supported by the following observations. First, *H. longicornis* is the most abundant of the HYSV-positive ticks in the endemic regions, and widely distributed in the Asia-Pacific region, including China, Korea, Japan, Australia, the Pacific Islands, and New Zealand [34–38]. Second, four migratory wild fowl species that are regular hosts of *H. longicornis* (*Zoothera aurea*, *Turdus hortulorum*, *Halcyon coromanda*, and *Pitta nympha*) are known to breed and migrate between China, South Korea, and Japan [39–42]. Third, the tick distribution matches the migratory bird flyways between China, South Korea, and Japan [43*,44]. Jeju Island, Korea, lying on the East Asia/Australasia Flyway, is home to one of the largest populations of migratory birds in Korea, and the place where 10 of 36 HYSV cases were identified [33**].

Taking the above observations together, a tick and migratory bird model for HYSV transmission is proposed here (Figure 1). Human infections by ticks from domestic animals and agricultural activities have been extensively recorded. HYSV has been isolated from ticks of cattle, goats, dogs and rats. Virus RNA has been detected in ticks from pigs and tea bushes, and domestic animals in endemic regions have high seroprevalence. The ticks of *H. longicornis* and *I. nipponensis* have been frequently found on wild boar, roe deer, raccoon dogs, Siberian weasels, Asian badgers, and leopard cats [24,45]. However, while the infection of humans by virus positive ticks from wildlife is very likely, more evidence is needed for the case of this specific pathogen. The following questions remain to be answered: where did the virus positive ticks from domestic animals originate, and how do migratory birds distribute virus positive ticks? This model needs to be further tested, but it may provide new strategies for the prevention and control of HYSV.

Conclusion

In 2009, a cluster of patients, from the Huaiyangshan mountain areas on the border of Hubei and Henan provinces of China, who had high fever, leucopenia, thrombocytopenia and elevation of liver transaminase levels, was first noticed at the Union Hospital of Tongji Medical College. A novel tick-borne Bunyavirus was identified as the causative agent. It was first named as HYSV, and later named as SFTSV. The disease is a hemorrhagic fever-like disease, and is widely distributed in China, South Korea and Japan. The clinical presentations of severe fever and thrombocytopenia are consistent with many bacterial or

virus infections. Of the five species of ticks positive for Huaiyangshan virus RNA, *H. longicornis* was the most abundant in endemic regions, and is considered to be the major vector. Its main hosts in endemic regions include cattle, goats, dogs, chickens, and rats, which were widely infected with HYSV as evidenced by various levels of seroprevalence. The *H. longicornis* ticks were hypothesized to play a central role in transmitting the virus, at least in part, by being transported over long distances to Korea and Japan by migratory birds. On the basis of the above observations, a tick and migratory bird model for HYSV transmission was proposed.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (81290340, 81290344-5 and 812111251).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Zhang YZ, Zhou DJ, Xiong Y, Chen XP, He YW, Sun Q, Yu B, Li J, Dai YA, Tian JH, Qin XC, Jin D, Cui Z, Luo XL, Li W, Lu S, Wang W, Peng JS, Guo WP, Li MH, Li ZJ, Zhang S, Chen C, Wang Y, de Jong MD, Xu J: **Hemorrhagic fever caused by a novel tick-borne Bunyavirus in Huaiyangshan, China.** *Zhonghua Liu Xing Bing Xue Za Zhi* 2011, **32**:209-220.
 2. Yu XJ, Liang MF, Zhang SY, Liu Y, Li JD, Sun YL, Zhang L, Zhang QF, Popov VL, Li C, Qu J, Li Q, Zhang YP, Hai R, Wu W, Wang Q, Zhan FX, Wang XJ, Kan B, Wang SW, Wan KL, Jing HQ, Lu JX, Yin WW, Zhou H, Guan XH, Liu JF, Bi ZQ, Liu GH, Ren J, Wang H, Zhao Z, Song JD, He JR, Wan T, Zhang JS, Fu XP, Sun LN, Dong XP, Feng ZJ, Yang WZ, Hong T, Zhang Y, Walker DH, Wang Y, Li DX: **Fever with thrombocytopenia associated with a novel Bunyavirus in China.** *N Engl J Med* 2011, **364**:1523-1532.
 3. Qu J, Zhang S, Li JD, Li DX: **Epidemic situation and surveillance on SFTS in China, 2011–2012.** *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 2013, **27**:257-259.
 4. Liu K, Zhou H, Sun RX, Yao HW, Li Y, Wang LP, Mu D, Li XL, Yang Y, Gray GC, Cui N, Yin WW, Fang LQ, Yu HJ, Cao WC: **A national assessment of the epidemiology of severe fever with thrombocytopenia syndrome, China.** *Sci Rep* 2015, **5**:9679.
 5. Kim KH, Yi J, Kim G, Choi SJ, Jun KI, Kim NH, Choe PG, Kim NJ, Lee JK, Oh MD: **Severe fever with thrombocytopenia syndrome, South Korea, 2012.** *Emerg Infect Dis* 2013, **19**:1892-1894.
 6. Takahashi T, Maeda K, Suzuki T, Ishido A, Shigeoka T, Tominaga T, Kamei T, Honda M, Ninomiya D, Sakai T, Senba T, Kaneyuki S, Sakaguchi S, Satoh A, Hosokawa T, Kawabe Y, Kurihara S, Izumikawa K, Kohno S, Azuma T, Suemori K, Yasukawa M, Mizutani T, Omatsu T, Katayama Y, Miyahara M, Ijuin M, Doi K, Okuda M, Umei K, Saito T, Fukushima K, Nakajima K, Yoshikawa T, Tani H, Fukushi S, Fukuma A, Ogata M, Shimajima M, Nakajima N, Nagata N, Katano H, Fukumoto H, Sato Y, Hasegawa H, Yamagishi T, Oishi K, Kurane I, Morikawa S, Saijo M: **The first identification and retrospective study of severe fever with thrombocytopenia syndrome in Japan.** *J Infect Dis* 2014, **209**:816-827.
 7. Xing Z, Schefers J, Schwabenlander M, Jiao Y, Liang M, Qi X, Li C, Goyal S, Cardona CJ, Wu X, Zhang Z, Li D, Collins J, Murtaugh MP: **Novel Bunyavirus in domestic and captive farmed animals, Minnesota, USA.** *Emerg Infect Dis* 2013, **19**:1487-1489.
 8. Bao CJ, Guo XL, Qi X, Hu JL, Zhou MH, Varma JK, Cui LB, Yang HT, Jiao YJ, Klena JD, Li LX, Tao WY, Li X, Chen Y, Zhu Z, Xu K, Shen AH, Wu T, Peng HY, Li ZF, Shan J, Shi ZY, Wang H: **A**

- family cluster of infections by a newly recognized Bunyavirus in eastern China, 2007: further evidence of person-to-person transmission. *Clin Infect Dis* 2011, **53**:1208-1214.
9. Wu Y, Gao GF: **Severe fever with thrombocytopenia syndrome virus expands its borders.** *Emerg Microbes Infect* 2013, **2**:e36.
 10. Yu B, Wang WY, Tian JH, Kong DY, Quan YX, Dai YA, Wang QF, Liu TY, Hu Q, He YW: **Epidemiological study on data involving 61 hospitalized cases with Huaiyangshan hemorrhagic fever in Wuhan.** *Zhonghua Liu Xing Bing Xue Za Zhi* 2012, **33**:124-125.
 11. Li W, He YW: **Infection with a novel virus causes hemorrhagic fever in China.** *Int J Infect Dis* 2013, **17**:e556-e561.
The infection caused by the newly discovered tick-borne Bunyavirus displayed diverse clinical symptoms and laboratory findings, as reported by the doctors who received and treated the first group of patients in China.
 12. Zhang L, Liu Y, Ni D, Li Q, Yu Y, Yu XJ, Wan K, Li D, Liang G, Jiang X, Jing H, Run J, Luan M, Fu X, Zhang J, Yang W, Wang Y, Dumler JS, Feng Z, Ren J, Xu J: **Nosocomial transmission of human granulocytic anaplasmosis in China.** *JAMA* 2008, **300**:2263-2270.
 13. Xu B, Liu L, Huang X, Ma H, Zhang Y, Du Y, Wang P, Tang X, Wang H, Kang K, Zhang S, Zhao G, Wu W, Yang Y, Chen H, Mu F, Chen W: **Metagenomic analysis of fever, thrombocytopenia and leukopenia syndrome (FTLS) in Henan Province, China: discovery of a new Bunyavirus.** *PLoS Pathog* 2011, **7**:e1002369.
 14. Zhang YZ, He YW, Dai YA, Xiong Y, Zheng H, Zhou DJ, Li J, Sun Q, Luo XL, Cheng YL, Qin XC, Tian JH, Chen XP, Yu B, Jin D, Guo WP, Li W, Wang W, Peng JS, Zhang GB, Zhang S, Chen XM, Wang Y, Li MH, Li Z, Lu S, Ye C, de Jong MD, Xu J: **Hemorrhagic fever caused by a novel Bunyavirus in China: pathogenesis and correlates of fatal outcome.** *Clin Infect Dis* 2012, **54**:527-533.
 15. Hiraki T, Yoshimitsu M, Suzuki T, Goto Y, Higashi M, Yokoyama S, Tabuchi T, Futatsuki T, Nakamura K, Hasegawa H, Saijo M, Kakihana Y, Arima N, Yonezawa S: **Two autopsy cases of severe fever with thrombocytopenia syndrome (SFTS) in Japan: a pathognomonic histological feature and unique complication of SFTS.** *Pathol Int* 2014, **64**:569-575.
 16. Zhang YZ, Zhou DJ, Qin XC, Tian JH, Xiong Y, Wang JB, Chen XP, Gao DY, He YW, Jin D, Sun Q, Guo WP, Wang W, Yu B, Li J, Dai YA, Li W, Peng JS, Zhang GB, Zhang S, Chen XM, Wang Y, Li MH, Lu X, Ye C, de Jong MD, Xu J: **The ecology, genetic diversity, and phylogeny of Huaiyangshan virus in China.** *J Virol* 2012, **86**:2864-2868.
 17. Huang X, Liu L, Du Y, Wu W, Wang H, Su J, Tang X, Liu Q, Yang Y, Jiang Y, Chen W, Xu B: **The evolutionary history and spatiotemporal dynamics of the fever, thrombocytopenia and leukocytopenia syndrome virus (FTLSV) in China.** *PLoS Negl Trop Dis* 2014, **8**:e3237.
Phylogenetic analyses indicated that the novel tick-borne Bunyavirus likely originated somewhere in Huaiyangshan circa 1790, and subsequently spread to other parts of China.
 18. Lam TT, Liu W, Bowden TA, Cui N, Zhuang L, Liu K, Zhang YY, Cao WC, Pybus OG: **Evolutionary and molecular analysis of the emergent severe fever with thrombocytopenia syndrome virus.** *Epidemics* 2013, **5**:1-10.
 19. Yoshikawa T, Shimajima M, Fukushi S, Tani H, Fukuma A, Taniguchi S, Singh H, Suda Y, Shirabe K, Toda S, Shimazu Y, Nomachi T, Gokuden M, Morimitsu T, Ando K, Yoshikawa A, Kan M, Uramoto M, Osako H, Kida K, Takimoto H, Kitamoto H, Terasoma F, Honda A, Maeda K, Takahashi T, Yamagishi T, Oishi K, Morikawa S, Saijo M: **Phylogenetic and geographic relationships of severe fever with thrombocytopenia syndrome virus in China, South Korea, and Japan.** *J Infect Dis* 2015, **212**:889-898.
Phylogenetic analyses indicated that, on rare occasions, the novel tick-borne Bunyavirus was transmitted between countries.
 20. He CQ, Ding NZ: **Discovery of severe fever with thrombocytopenia syndrome Bunyavirus strains originating from intragenic recombination.** *J Virol* 2012, **86**:12426-12430.
 21. Ding NZ, Luo ZF, Niu DD, Ji W, Kang XH, Cai SS, Xu DS, Wang QW, He CQ: **Identification of two severe fever with thrombocytopenia syndrome virus strains originating from reassortment.** *Virus Res* 2013, **178**:543-546.
 22. Wang S, Li J, Niu G, Wang X, Ding S, Jiang X, Li C, Zhang Q, Liang M, Bi Z, Li D: **SFTS virus in ticks in an endemic area of China.** *Am J Trop Med Hyg* 2015, **92**:684-689.
The novel tick-borne Bunyavirus may be transmitted via the transovarial route, as evidenced by detection of viral RNA in *H. longicornis* at all developmental stages.
 23. Meng K, Sun W, Cheng Z, Guo H, Liu J, Chai T: **First detection of severe fever with thrombocytopenia syndrome virus in the tick species *Haemaphysalis concinna* in Shandong Province, China.** *Parasitol Res* 2015.
 24. Park SW, Song BG, Shin EH, Yun SM, Han MG, Park MY, Park C, Ryou J: **Prevalence of severe fever with thrombocytopenia syndrome virus in *Haemaphysalis longicornis* ticks in South Korea.** *Ticks Tick Borne Dis* 2014, **5**:975-977.
 25. Yun SM, Lee WG, Ryou J, Yang SC, Park SW, Roh JY, Lee YJ, Park C, Han MG: **Severe fever with thrombocytopenia syndrome virus in ticks collected from humans, South Korea, 2013.** *Emerg Infect Dis* 2014, **20**:1358-1361.
Of the novel tick-borne Bunyavirus positive ticks collected from humans in South Korea, the *H. longicornis* ticks was the most abundant, followed by *Amblyomma testudinarium*, and *Ixodes nipponensis* ticks.
 26. Chang MS, Woo JH: **Severe fever with thrombocytopenia syndrome: tick-mediated viral disease.** *J Korean Med Sci* 2013, **28**:795-796.
 27. Ding S, Yin H, Xu X, Liu G, Jiang S, Wang W, Han X, Liu J, Niu G, Zhang X, Yu XJ, Wang X: **A cross-sectional survey of severe fever with thrombocytopenia syndrome virus infection of domestic animals in Laizhou City, Shandong Province, China.** *Jpn J Infect Dis* 2014, **67**:1-4.
 28. Niu G, Li J, Liang M, Jiang X, Jiang M, Yin H, Wang Z, Li C, Zhang Q, Jin C, Wang X, Ding S, Xing Z, Wang S, Bi Z, Li D: **Severe fever with thrombocytopenia syndrome virus among domesticated animals, China.** *Emerg Infect Dis* 2013, **19**:756-763.
 29. Li Z, Hu J, Bao C, Li P, Qi X, Qin Y, Wang S, Tan Z, Zhu Y, Tang F, Zhou M: **Seroprevalence of antibodies against SFTS virus infection in farmers and animals, Jiangsu, China.** *J Clin Virol* 2014, **60**:185-189.
Using a double-antigen sandwich ELISA, the team found that the seroprevalence for the novel tick-borne Bunyavirus in domestic animal species from endemic regions were: goats (66.8%), cattle (28.2%), dogs (7.4%), pigs (4.7%), chickens (1.2%), geese (1.7%), rodents (4.4%) and hedgehogs (2.7%).
 30. Xing Z, Scheifers J, Schwabenlander M, Jiao Y, Liang M, Qi X, Li C, Goyal S, Cardona CJ, Wu X, Zhang Z, Li D, Collins J, Murtaugh MP: **Novel Bunyavirus in domestic and captive farmed animals, Minnesota, USA.** *Emerg Infect Dis* 2013, **19**:1487-1489.
 31. Riemersma KK, Komar N: **Heartland virus neutralizing antibodies in vertebrate wildlife, United States, 2009-2014.** *Emerg Infect Dis* 2015, **21**:1830-1833.
 32. Ni H, Yang F, Li Y, Liu W, Jiao S, Li Z, Yi B, Chen Y, Hou X, Hu F, Ding Y, Bian G, Du Y, Xu G, Cao G: ***Apodemus agrarius* is a potential natural host of severe fever with thrombocytopenia syndrome (SFTS)-causing novel Bunyavirus.** *J Clin Virol* 2015, **71**:82-88.
The RNA of the novel tick-borne Bunyavirus was detected in two of eight *Apodemus agrarius*, from which two strains were isolated. The genome sequences of isolates from patients and from *A. agrarius* had high levels of sequence identity.
 33. Yun Y, Heo ST, Kim G, Hewson R, Kim H, Park D, Cho NH, Oh WS, Ryu SY, Kwon KT, Medlock JM, Lee KH: **Phylogenetic analysis of severe fever with thrombocytopenia syndrome virus in South Korea and migratory bird routes between China, South Korea, and Japan.** *Am J Trop Med Hyg* 2015, **93**:468-474.
The authors hypothesize that the novel tick-borne Bunyaviruses in South Korea share common phylogenetic origins with samples from China and Japan, and that migratory birds, well-known carriers of the tick *H. longicornis*, are a potential source of HYSV transmission across countries. Thus, migratory birds play an important role in dispersing *H. longicornis*-borne viruses.

34. Zheng H, Yu Z, Zhou L, Yang X, Liu J: **Seasonal abundance and activity of the hard tick *Haemaphysalis longicornis* (Acari: Ixodidae) in North China.** *Exp Appl Acarol* 2012, **56**:133-141.
 35. Chen Z, Yang X, Bu F, Yang X, Yang X, Liu J: **Ticks (Acari: Ixodoidea: Argasidae, Ixodidae) of China.** *Exp Appl Acarol* 2010, **51**:393-404.
 36. Chen Z, Liu Q, Liu JQ, Xu BL, Lv S, Xia S, Zhou XN: **Tick-borne pathogens and associated co-infections in ticks collected from domestic animals in central China.** *Parasit Vectors* 2014, **7**:237.
 37. Chong ST, Kim HC, Lee IY, Kollars TM Jr, Sancho AR, Sames WJ, Chae JS, Klein TA: **Seasonal distribution of ticks in four habitats near the demilitarized zone, Gyeonggi-do (Province), Republic of Korea.** *Korean J Parasitol* 2013, **51**:319-325.
 38. Heath A: **Biology, ecology and distribution of the tick, *Haemaphysalis longicornis* Neumann (Acari: Ixodidae) in New Zealand.** *N Z Vet J* 2015:1-11.
 39. Ishiguro F, Takada N, Masuzawa T, Fukui T: **Prevalence of Lyme disease *Borrelia* spp. in ticks from migratory birds on the Japanese mainland.** *Appl Environ Microbiol* 2000, **66**:982-986.
 40. Trovao NS, Suchard MA, Baele G, Gilbert M, Lemey P: **Bayesian inference reveals host-specific contributions to the epidemic expansion of Influenza A H5N1.** *Mol Biol Evol* 2015.
 41. Lee DH, Torchetti MK, Winker K, Ip HS, Song CS, Swayne DE: **Intercontinental spread of Asian-Origin H5N8 to North America through Beringia by migratory birds.** *J Virol* 2015, **89**:6521-6524.
 42. Kang HM, Batchuluun D, Kim MC, Choi JG, Erdene-Ochir TO, Paek MR, Sugir T, Sodnomdarjaa R, Kwon JH, Lee YJ: **Genetic analyses of H5N1 avian influenza virus in Mongolia, 2009 and its relationship with those of eastern Asia.** *Vet Microbiol* 2011, **147**:170-175.
 43. Kim BJ, Kim H, Won S, Kim HC, Chong ST, Klein TA, Kim KG, • Seo HY, Chae JS: **Ticks collected from wild and domestic animals and natural habitats in the Republic of Korea.** *Korean J Parasitol* 2014, **52**:281-285.
- Authors studied the distribution and density of ticks in 35 animals from five provinces and three cities in South Korea in 2012. They found that *Haemaphysalis longicornis* ticks were the most abundant.
44. Choi CY, Kang CW, Kim EM, Lee S, Moon KH, Oh MR, Yamauchi T, Yun YM: **Ticks collected from migratory birds, including a new record of *Haemaphysalis formosensis*, on Jeju Island, Korea.** *Exp Appl Acarol* 2014, **62**:557-566.
 45. Kim HC, Han SH, Chong ST, Klein TA, Choi CY, Nam HY, Chae HY, Lee H, Ko S, Kang JG, Chae JS: **Ticks collected from selected mammalian hosts surveyed in the Republic of Korea during 2008-2009.** *Korean J Parasitol* 2011, **49**:331-335.